



small dog

Flea Adulticide/Larvicide/Ovicide and Lice Adulticide

For Dogs 7 weeks and older weighing up to 4.5 kg

Once-A-Month Topical Flea and Lice Treatment Liquid

READ THE LABEL AND INSERT BEFORE USING FOR USE ONLY ON DOGS Do not use on other animals KEEP OUT OF REACH OF CHILDREN

WARNING POISON



EYE IRRITANT DOMESTIC

GUARANTEE: Imidacloprid 9.1 % and Pyriproxyfen 0.46 % REG. NO. 27581 P.C.P. ACT CONTENTS: One to Six 0.4 mL Tubes

Advantage® II small dog provides dual protection against fleas and lice. Advantage® II small dog kills 98–100 % of the existing fleas on dogs within 1 day after treatment. Protection against further flea infestation can last up to 4 weeks. Retreatment at monthly intervals is required if reinfestation is continuous or if flea control is desired. If reinfestation becomes necessary earlier than 4 weeks, do not retreat more than once weekly. Fleas, eggs, and larvae are killed following contact with skin debris from an Advantage® II small dog treated pet. Advantage® II small dog provides multi-stage flea control effectively breaking all flea life-cycle stages for quick and lasting control of flea populations. Advantage® II small dog treats lice infestations on dogs.

Do not shampoo or immerse the animal in water immediately after treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days after treatment. Advantage® II small dog retains its ovicidal and larvicidal efficacy up to 4 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be shortened by repeated shampooing, water immersion or exposure to rain. Reapplication may be required after such activities.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage® II small dog rapidly kills fleas and may reduce the incidence of this condition. For effective flea and lice control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. Do not get in eyes. Avoid contact with skin or clothing. Wear rubber gloves during application of this product. After handling or applying, wash hands (or any other skin that came in contact with the product), with soap and water. Avoid contact with pets until dry.

PETS: Do not use on animals under 7 weeks of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately. Call 1 888-663-5326 (1 888-NO-FLEA6) for assistance regarding Advantage® II small dog use.

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION:



Treat the patient symptomatically.

DISPOSAL:


Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose in household garbage.

DIRECTIONS FOR USE:

Do NOT administer by mouth. For dogs 7 weeks and older weighing up to 4.5 kg. This product is packaged in single dosage tubes. This package contains one to six monthly 0.4 mL dosages. Treat your dog monthly in the manner described below. Do not apply a dosage higher than one 0.4 mL tube at one time. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet. Retreatment at monthly intervals is required if reinfestation is continuous or if flea control is desired. If reinfestation becomes necessary earlier than 4 weeks, do not retreat more than once weekly. Use only on dogs. Do not use on other animals. Do not use on puppies under 7 weeks of age.



Remove one applicator from the package. Hold applicator tube in upright position. Pull cap off tube. Turn cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube.



The dog should be standing for easy application. Part the hair on the dog's back between the shoulder blades until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin. Do not get this product in your pet's eyes or administer by mouth. Until Advantage® II small dog dries, keep pet off furniture and away from hardwood floors. Discard the empty tube as described in "DISPOSAL".

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE © TM see www.bayer.ca/tm-mc

Bayer Inc. 2920 Matheson Blvd East Mississauga, Ontario L4W 5R6 1 888 663-5326



87070093

Table with product details: PM no., CR no., Format, Font size, Drawing no., Colours, Operator ID, Version no., Date, and contact information for questions.



chien de petite taille

Adulticide/larvicide/ovicide contre les puces et adulticide contre les poux
Pour chiens de 7 semaines et plus, pesant 4,5 kg ou moins
Traitement topique mensuel contre les puces et les poux
Liquide

LIRE L'ÉTIQUETTE ET LE FEUILLET D'EMBALLAGE AVANT L'EMPLOI
POUR USAGE SUR LES CHIENS SEULEMENT
Ne pas utiliser sur d'autres animaux
GARDER HORS DE LA PORTÉE DES ENFANTS

AVERTISSEMENT



POISON

IRRITANT POUR LES YEUX
USAGE DOMESTIQUE

GARANTIE : Imidaclopride à 9,1 %, pyriproxyfène à 0,46 %
N° D'HOM. 27581 L. P. A.
CONTENU : Un à six tubes de 0,4 mL

Advantage® II chien de petite taille protège contre les puces et les poux. Il tue, en deçà d'une journée du traitement, de 98 à 100 % des puces ayant infesté les chiens. Le produit peut protéger l'animal contre une réinfestation pour une période allant jusqu'à quatre semaines. Un traitement mensuel est nécessaire si l'animal est continuellement réinfesté ou si une action préventive contre les puces est désirée.

Ne pas faire de shampooing ni immerger l'animal dans l'eau immédiatement après le traitement. L'animal peut recevoir un shampooing, être immergé dans l'eau ou être exposé à la pluie trois jours après le traitement.

L'activité alimentaire répétitive des puces infestant les animaux de compagnie peut provoquer chez ces derniers un trouble d'hypersensibilité de l'épiderme : la dermatite allergique aux piqûres de puces (DAPP). Le traitement avec Advantage® II chien de petite taille tue rapidement les puces et diminue l'incidence de cette dermatite.

PRÉCAUTIONS :

ÉTRES HUMAINS : GARDER HORS DE LA PORTÉE DES ENFANTS. Nocif si ingéré ou si absorbé par la peau. Cause de l'irritation oculaire. Ne pas mettre dans les yeux.

ANIMAUX DE COMPAGNIE : Ne pas utiliser sur des animaux âgés de moins de 7 semaines. Faire attention à ne pas exposer les autres animaux qui peuvent se trouver à proximité de l'animal traité.

PREMIERS SOINS :

ÉTRES HUMAINS : EN CAS DE CONTACT AVEC LES YEUX, garder les paupières écartées et rincer doucement et lentement avec de l'eau pendant 15 à 20 minutes.

EN CAS D'INGESTION, appeler immédiatement un centre antipoison ou un médecin pour obtenir des conseils sur le traitement.

EN CAS DE CONTACT AVEC LA PEAU OU LES VÊTEMENTS, enlever tous les vêtements contaminés. Rincer immédiatement la peau à grande eau pendant 15 à 20 minutes.

ANIMAUX DE COMPAGNIE : Si des signes de sensibilité apparaissent, baigner l'animal en ajoutant à l'eau un savon doux, puis le rincer avec beaucoup d'eau.

RENSEIGNEMENTS TOXICOLOGIQUES : Traiter le patient selon les symptômes.

ÉLIMINATION : Ne pas réutiliser le contenant vide. Envelopper le tube vide et le bouchon dans du papier journal et jeter avec les ordures ménagères.


MODE D'EMPLOI :

NE PAS administrer par voie orale. Pour chiens de 7 semaines et plus, pesant 4,5 kg ou moins. Ce produit se présente sous forme d'un tube applicateur à dose unique.

Retirer un applicateur de l'emballage. Tenir le tube en position verticale, ouverture vers le haut.

Retourner le bouchon et le placer en position inversée au bout du tube.

Tourner le bouchon afin de briser le sceau du tube, puis retirer le bouchon du tube.



Retirer un applicateur de l'emballage. Tenir le tube en position verticale, ouverture vers le haut. Retirer le bouchon du tube. Retourner le bouchon et le placer en position inversée au bout du tube.



Pour faciliter l'application, maintenir le chien debout. Écarter le pelage du dos du chien entre les omoplates jusqu'à ce que la peau soit visible.

GARDER LE FEUILLET D'EMBALLAGE POUR RÉFÉRENCE FUTURE.

© MC voir www.bayer.ca/tm-mc

Bayer Inc.
2920 Matheson Blvd East
Mississauga (Ontario)
L4W 5R6
1-888-663-5326



Table with product information, contact details, and a list of names for questions.